

Concurrence Discrepancy at Level by Forced Replacement.

Nature of the Problem: determinating parameter ϑ in the probability distribution function $f(x \mid \vartheta)$ as unknown as a File. (Access Review or *Illustrée as this Paper*) Belonging to an Interval Ω in $\mathbb{R}$. (observed values in sample of secondary effects). We estimate ϑ . Comparative Estimator in Discrepancy and relation to this document. An objective is for me is to proceed and document. This is as Digital Experiment is with Content in Bayesian Estimation. The Completion is an Activity supported by an Assistant in Parallelism. See Local Areas Networks and Priorities specific to Clozaril as an Image. See Indemnization form Allan Memorial. (as an Experiment). The Access Review: from Sample (s_i, y_i) , s_i a suite of events: Contingent: and $f: x_i \rightarrow y_i$ at $f(g(a_{ij})) = f \circ g_i(x_i)$ as Probability distribution function: Gain from a Randomized Event and Access Reviews. (No variance in $x_i \rightarrow y_i$ is by Half Line as $x_i \rightarrow y_i$ (data is a probability distribution function as frequency) where $x_i < y_i$ or $y_i < x_i$, One wants to find as a particular ordered sequence of n items with discrepancy (defective or non defective). *Suivi* in the Medium as Access Review.

Passage and Path for Level.

$(X = X_1 + X_2 \dots + X_n$ where X_i and X has a distribution with parameter n and p).

We assign $\Pr(X_i) = C_{n,x_i} p^{x_i} (1-p)^{n-x_i}$ and observe that $\Pr(X) = \Pr(\sum X_i) = np$ and $\sigma = npq$.

In this problem we have a set of n , with proportion p , and $x_i \in [0; n] \cap \mathbb{N}$ and $x_i = X_i$. Another way to set the quantities is to say the step is p , time n and availability x_i .

Waiting.

There the time is $[0; t]$, the number of events x_i in time, and close to λ in all.
 $e^x = 1 + x + \frac{x^2}{2!} + \dots$ and $e^\lambda e^{-\lambda} = 1 = \frac{\lambda e^{-\lambda}}{1!} + \dots + \frac{\lambda^n e^{-\lambda}}{n!} + \dots$

$$E\left(\frac{1}{e^\lambda} \sum p(x)\right) = E\left(\frac{e^x}{e^\lambda}\right) = 1$$

The number if x_i in event E_i in t , set in $i \leq n$ (recall $x_i = i$), giving $p(x) = \frac{\lambda^{x_i}}{x_i!} e^{-\lambda}$ that must be skewed to the left, as for a large threshold $\frac{\lambda^{x_i}}{x_i!}$ will become very small.

Experience and Concurrence.

$1 + x + x^2 + \dots = \frac{1}{1-x}$ as x and $x^k \in (-1; +1)$ for $k \in \mathbb{N}$, and $1 : (1-x)$, as all : false.

$$f(x) = \left\{ \begin{array}{l} \frac{1}{k} \text{ for } x = 1, \dots, k \\ 0 \end{array} \right\} \rightarrow \text{Uniform distribution on the integers } 1, \dots, k, \text{ No}$$

Psychosomatic feel. (Elastic). *Que la vie aie l'air d'un film parfait.*

$1 + x^2 + x^3 + \dots = \frac{1}{1-x} \forall x \in [-1, 1]$, as **One Variable Calculus Job : Order** (see **Paper on Loops Ordering for Sort for *Illustrées Access Reviews*** as Experience). Also by Riemann's Sums as *Illustrées*. (an Expansion at Groups in Snagov). The Lebesgues Sustainable Measure is as *Illustrées* by Surjectivity with Margin and Domains in $[c, d]$ as Equidistributed Sequence (as *Illustrée*) (modification by minimality and Skeweness to Left).

Measure as Constraint and Access Reviews: columns as $a_j = f_i$ as Stralen's of Burger and common at Right Top Corner as Rest with Deals: with Expanded Terms: Deals:

$$\sum_{i=1}^{\infty} \frac{1}{n(n+1)} = \frac{1}{1 \cdot 2} + \frac{1}{2 \cdot 3} + \dots (\text{Entropy}) \sum_{i=1}^{\infty} \frac{1}{n(n+1)} = \sum_{i=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n+1} \right) : \text{Telescoping by Snagov.}$$

The constraint and Movement Cone is as Equidistant Sequences (Equidistributed) and with Chernikova's and Vertices as Equidistant.

Loops and Ordering for Sort and Discrepancy.

Object of the **Premisse**: reasoning by recurrence: Bankok and Bourbaki: Good for 0 and 1, and for $n \in \mathbb{N}$, then $n + 1$ valid : a property for all $\mathbb{N}$. Defined as Ambiance. A family v_i is free if v_i , are linearly independent: as linear programming by orthonormalization of v_i , by Gram Schmidt. The **Loop** is by GraphTree where nodes ridges: have as extremities same node as networks. **The Loop is as: portion of code as many times in a row: until a condition drives you out from Loop.** Safeguard. The condition may not function some times and goes to ∞ without interruptions: as mono task: where multi tasks comes to first tasks at beginning of iteration.

While is defined: Pré-Condition as Condition of Stop: where $\mathbb{N}$ iteration and Post-Condition at **Do or Until** as branched Loop making a List as a Casino.

The Tree is by: *arête ayant pour extrémité même Sommet: Compteurs (associé a une variable incrementée de façon variable dans la boucle et Iterateurs pour différents types pour ces Boucles.*

Control Structure and Recuperation: Rosenthal and Audience where teaching is tom a Group over time. Branding and Parallelism: as Home Detox: Insurance and Buying Insurance. A Clinical Practitioner and Parallelism of Floor is for Relaxation $O(n)$ with Home Proof: Safety with Half Line Continuity: see Fitbit. The Identification as elimination of toxines eliminated from Room: Maintenance and MUHC and Tree Traversal: Access to Ambiance: as Home Domain, Method as $O(n)$, Kitchen Bathroom Bedroom LivingRoom LaundryRoom and Home Office: Ressources for Toxin Tests Sewage Ridge: Home Detox: Restoration of Health: $O(n)$ Exposure in $\mathbb{N}$ Tree Exposure for Health: with Range as no Inflammation: and DNA Risk: suffering and Quality of Life: triggering In and Out (Loop for Protection: Onto Subjective and defective $O(n)$).

Reduced Kidney Function: Air Surface of Home as Application and Walls: determining Path. Detox as Map: from Generic Graph *Domain* $\rightarrow$ *Range* as Tract Groin at MUHC as *Root* $\sqrt{x}$ and $\text{sgn}(x) = \frac{x}{|x|}$. Number Theory for *Les Étapes*: Convexity of f as Straight Line or Circle: As such Chernikova: an example $f(x) = 2x - 3$: Slope 2, Intercept -3, $g(x) = \sqrt{16 - x^2}$, $\sqrt{x+1} = f(x)$, $g(x) = \frac{x-1}{x}$ at $f+g, fg, \frac{f}{g}$. End Growth and Finite Increments as f and g and $|f'| < |g'|$. The tree at Welfare at Destination $\Delta f = \Delta g$ as $|f(b) - f(a)| \leq |g(b) - g(a)|$. The Luxury is defined as Circularity and $O(n)$ is by the Wall: McGill.


Path Increment (Diet and Removing) : Onto Cells and Applications: discerning f and g . Outer Loop: metals pesticides fungi bacteria and Inner liver kidneys lungs lymph tract sweat locating toxines. Tree for Well Being as Network: toys and body for consistent exposure by hidden toxines and blue print of toxines: Energy and Removing.

Path Method as Determination: Maintenance Relaxation: Loops as sponges brushes vacuum All Rooms Air Purifiers brooms Dustpans Brushes


From The Right of Principality: π_i and $O(n_i)$ where Super Toxins as $x \in \partial G$: Loop as asthma, headaches, inflammation and fatigue: Kidneys toxines Skin Care: No Synthetic and mineral and Plants. π_i as Home Detox by Love and Social Order: Soins Santé and Social

Services for Work and Policy in $O(n)$ to Resume: emotional Intelligence: see Ticket and Catastrophy. member@npqmag.org in Book of Home Detox.

Juridic Right, Home and Loops: happens that Inner Loops run many times inside some Outer Loops: and there is Interchange: two Iteration Variables used by nested or Outer Loops. The Ellipsoid and Higher Social Stand at Snagov. The Limit behaviour als Ordnung $f(x) = O(g(x))$, $x \rightarrow \infty, M > 0$. Perpendicularity with some $x_0 = 5, 0 \leq f(x) \leq Mg(x), x > x_0$. Here $f(x) - Mg(x) \leq 0$ as Run Time or Space requirement. Recall that the Growth Rate is by f as a Valued Function (forwarding f) and g_i as a Comaprison (corrector g_i). We have Domain as Unbounded Set (Half Line): $f(x) = O(g(x))$. If $|f(x)|$ is at most a positive Constant multiple of $g(x)$, $\forall$ large x . If $\exists M > 0, \exists x_0$ such that $|f(x)| \leq Mg(x)$, as $x \geq x_0$. Therefore $|f(x)| - Mg(x) \leq 0$ and $f(x) = O(g(x))$.



In the Loop: we may have asymptotic Growth Rates as $g(x)$ from $f(x)$. (see Tuineag)
 One may determine a Pension Fund by $O(n)$, called Order of Function: upper Bound of Growth Rate of Forward as Evidence: with Majorant as Subset of Preordered of K preordered : Majorant $\in$ Subset and Minorant as Subset of Preordered of K preordered : Minorant $\in$ Subset. Majorized or Minorated as bounded form above and below with Left Continuity:




Problem: f and g_i , defined on some **unbounded** Subset of $x \in \mathbb{R}^+$. Here unbounded as No Boundary (No finite Major or Minor and no Measure). A circle is boundary less but a Half Plane is as Left Continuity: Not necessarily a Closed Set. $f(x) = x^2 + 1$, $g(x) = x^2 - 1$. Constrained on $S \subset \mathbb{R}^2$ and is closed. Well defined on $\mathbb{R}$ but Unbounded.

$|f(x)| \leq Mg(x)$, $\forall x \geq x_0$ as $x \rightarrow \infty$ and no $x = a$ but $x \rightarrow a$. Here $0 \leq |x - a| < \delta$, $\forall x$.
 $|f(x)| \leq Mg(x)$ with $g(x)$ strictly positive for all x close to a . See

$$\limsup_{x \rightarrow a} \frac{|f(x)|}{g(x)} < \infty$$

The limit point as Critical and Singular $a \in \mathbb{R}$, (whether $a = \infty$ or not) is a cluster point of $D(f)$ and $R(f)$. There is an Accumulation Point as Critical and Singular. $\forall f \& g$ of some unbounded Set as Accumulation Point of $i \in \mathbb{N}$, $\forall x \in \text{Range } \mathbb{R}^+$ as $f(x) = O(g(x))$, if there exists M , with x_0 such that $f(i) \leq Mg(i)$, $\forall i \geq x_0$. Example: for large x , $f(x) = 6x^4 - 2x^3 + 5$ as $x \rightarrow \infty$. The Highest Growth Rate is $6x^4$ as $|f(x)| \leq Mg(x)$. $f(x) = O(x^4)$.
 $|6x^4 - 2x^3 + 5| \leq Mx^4$. How closely a finite series approximates a given function? g_i is a simple function omitting Constant factors and lower order terms. Infinite Assymthoties with different limits for the function argument: The number of Steps of Loop is the $n^{\text{biggerPower}} = n^x$ dominating. The **coefficient irrelevant as \$ Support**: Infinitesimal Asymptotic: Error Term (Approximation). Significant terms: Taylor's Polynomial or Residual with Size of Reminders



$\sin x \approx x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!}$ on Support $-\pi \leq x \leq \pi$. Infinite Asymptotic: $n!, 2^n, n^2, n \log n, n$. Infinitesimal Asymptotic: Residual in Gandia: $O(n^c)$ & $O(c^n)$. if $c \geq 1$ and $O(c^n)$ a Super Polynomial.

$$O(\log n) = O(\log n^c).$$

$f_1 f_2 = O(g_1 g_2)$, $\forall f_i = O(g_i)$ and $f \cdot O(g) = O(fg)$, and by $f_1 + f_2$ as $O(\max g_1 g_2)$ a convex cone as Chernikova's Cone or other convex cone.

Multiple Variables and Help with Vectors and Pain $[x_i] = \vec{x}$. Let $f(x_i)$ is $O(g(x_i))$, $[x_i] \rightarrow \infty$ iff $\exists M, c \geq 0$ such that $|f([x_i])| \leq c |g([x_i])|$, $\forall [x_i] \geq M, i \in \mathbb{N}, \|[x_i]\|_\infty \geq M$ the Chebyshev Norm. Example: $f(n, m) = n^2 + m^3 + O(n + m)$ as $n, m \rightarrow \infty, \exists c, M$ such that $|f(n, m) - [n^2 + m^3]| \leq c|n + m|$, $m \geq M, n \geq M$. There are Multiple Variables as Home Detox Complexity or 2^{nd} dimension.

$$O(1)$$

$O(\log \log n)$:Interpolating Research with Key in Array ordered by i as x_i numerical value order.

$O(\log n)$:Balanced Tree Search: Binary and Half Interval Search: Position of Target Value in a sorted Array.

$$O((\log n)^c), c \geq 1, \text{ Matrix chain Ordering (isotope)}$$

$$O(n^c), 0 \leq c \leq 1 \text{ in Tree}$$

$$O(n) \text{ as linear}$$

$$O(n \log^* n) \text{ at } \log^*(n) = \begin{cases} 0 \\ 1 + \log^* \log(n) \end{cases}$$

$O(n \log n) = O(\log(n!))$ a Merge Sort at PeepShow (Divide and Conquer Algorithm)- at all neighbors.

$$O(n^2) \text{ Buble Insertion and Quick Sort}$$

$O(n^c)$ Tree adjoining Grammar: Parsing matching and Bipartite Graph: LU decomposition.

$$O(c^n) \text{ Salesman Problem: travelling Brute Force Search.}$$

The Salesman Problem and Territory: from List of Cities and Distances: find order in shortest possible visits of each City and come back to Origin.

The Peep Shop: Moving to Ethical Device: Accessible Story and Ticket by Catastrophy. Telling an actionable frame work for comprehensive ethical Risk at Organization: Data

Science and Étude du Point: at the Geek Squad Community: Transfer as **Merge Sort with Arab** $O(n \log^*(n))$ as **Cairo Rare and Algiers: $O(n)$ Artiste Peintre Illustration as Application with Left Right Joint and Catastrophy and Ticket as Maintenance as a Cone where Chernikova's Algorithm suits as Erdos Number where Maintenance comes as a Cone as Property of Hamams.** (Vertices and Loops). Cairo Rare. Equivoque as both ways map (same invertibility) and Univoque as One to One and Reverted, with Biunivoque as same map for inversion and One to One. Cartography.

Territory Surveillance and Health Costs. (presence of Shears as Spatial Turn)

Spatial Turn is a term used in conjunction with iCloud. It relates Origin with Border (as a polar coordinate specifying Work statistics)

Bordering is defined: **Feedback Return** definition is: $\exists f_i$ tangent to E , a bordered convex set as a closed Loop or closed Sequence, this closing E , and determining it. Open Loop is defined as an Open Set and the same for Closed.

Determinism is defined: **The Hahn Banach and Separation** theorem introduce a Work function at π_i at $i = k$. For these, $\exists \mathbb{P}$ a Sphere as given around an Origin, and $P \notin \mathbb{P}$, then $\exists \pi_k$ hyperplanes, with $P < \pi_k < \mathbb{P}$.

Dialectics and **Duality** are regularly introduced as:

$$\min_{\mathbb{P}}(P - \mathbb{P}) = \max_{K \text{ to } \mathbb{P}}(P_k - \pi_K(P)), \quad \forall \pi : P < \pi_K < \mathbb{P}$$

It is seen as: from **Shears we meet a Two Track Research Approach, and are the Work part of the Product.**

The **Hahn Banach Theorem and Work Extension** is seen as X a normed linear vectorspace, $M \subseteq X$, $\mathbb{D} \in M^*$, a bounded linear functional on M , $\exists \tilde{\mathbb{D}} \in X^*$ extending the bounded linear functional and $|\mathbb{D}| = |\tilde{\mathbb{D}}|$.

Ordering is defined: the Bubble Sort Algorithm, with the **Influence as Job Specification** where $A : (a_{ij})$ is a Role **Data Shift** $E_1 \otimes E_2$ and is:

$$\text{Projection}[E_2] \subset \text{Projection}[E_1] \text{ with } x \leq A^{-1}b \text{ an } \mathbf{Orangerai}$$

Othoring is defined: **Venue is by Augmented Reality**: $x_{.i}$ in $AX = B$ is a solution to $\min CX - d$ subject to $AX + B, X \geq 0$. We call $x_{.i}$ a feature of Venue. **Work is the dual of the Augmented Reality**: $\max B^T U - d$ subject to $A^T U \leq C^T$, where $u_{.i}$ is a feature of Work.

Repère is defined as Inflection Points among Critical Points.

Borderland is defined as: Fuzzy Logic with k nearest neighbor Algorithm.

Excessive firing on $x_i \rightarrow y_i$ is by a Scientific Border x_i to Over Determination y_i .

Network Border is defined as: The **stable** part of E provided by an inner product with: $a, b \in A \subset E$, with $a * b \in A \subset E$.

The **stable** part of set E with action $*$ of Ω on E , $(\alpha, x) \in \Omega \otimes A$, $\alpha * x \in A \subset E$. The **stable** part of f of $E \rightarrow E: P \subset E$, such that $f(P)$ of P by $f \subset P$. **Stationary point**: of an arc parametrized (I, f) of $\mathbb{R}^3$ of class C^k with $k \geq 2$ (not an ordinary point), $M \in \text{arc}$ such that $\frac{\partial \vec{M}}{\partial t} = \vec{f}'(t) = 0$. Here in this Border, Immigration by Walk is addressed.

Borderscape: imaginative frontier when using the Software.

Rebordering is defined as: **Commands in Optimal Time are close to Google Drive.**

$[x^i(t)]$ are Phase coordinates. $[u^i(t)]$ command coordinates. See $[x^i(t)] \in X$ the Phase Space, and the admissible Command $[u^i(t)]$ may lead to $[u^i] \in \mathbb{R}^r$, with the closed domain

of Command Space $U \subset \mathbb{R}^r$. The Objective of the definition: The Border and Colonialism and Russia and Spain lead to Ground x_i . Group Funding as y_i . **The Wealth Pivot is defined in a_{ij} in Economy as Purchase.** The Employee $x_i \rightarrow$ industrial relation y_i and is defined at Organization by Individual Business nad Gouvernement.

Representation is defined as Babel+ Productivity in I_i . (Daily Online Leader) The Ends Meet $x_i \rightarrow y_i$ is formal in x_i defined as: $\exists i$ from b_i as Presence in Calendar as Casual for transitivity. One has flexible Incentive and Added Value in Gandia. The Expats and Agency is by Support for Reporting. The objective is no Re Location to settle in UE as a Path forward to Companies in Gandia and UE. The argument is: Growth, Ressource, Dedication. Asset, Goals, ?employee, Investment $\leftrightarrow$ Ellipse and Traditional training, Lean and Digitization. Praxis is by $\leq b_i$ discrete as a symbol suite for partition. The x_0 is form the Cache as Establishment.

The Economic Forum is from Business Industry and Ends Meet, y_i, i , in Russian Foreign Policy as totally bounded. From Plaza Coffee Massage Store Bus Pivot a_{ij} , Bibliothèque and Math in Pivot.

Sustainability is by a Lagrangian: Vector Calculus $\rightarrow$ Indefinite Integral:
 $\exists path : \sigma : [a; b] \rightarrow \mathbb{R}^n$ differentiable $\exists \sigma'(t_0)$ as tangent defined as $\sigma(t_0) + \lambda \sigma'(t_0)$. For **Sustainability** we have the investment space set $i \rightarrow$ a totally bounded suite $\forall i \exists \leq B$. This is by creation of Concession. This Set as an investment space has $\mathbb{N} \rightarrow \ln|X_i| \in Market$.